



Recombinant Human IL-2 (aldesleukin)

Catalog #	EPT147
Expression Host	E.coli
DESCRIPTION	Recombinant Human Interleukin-2 is produced by our E.coli expression system and the target gene encoding Pro22-Thr153(Cys145Ser) is expressed.
Accession	P60568
Synonyms	Interleukin-2; IL-2; T-cell growth factor; TCGF; Aldesleukin
Mol Mass	15.5 KDa
AP Mol Mass	14 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test.
FORMULATION	Lyophilized from a 0.2 μ m filtered solution of 10mM Acetata-Na, 5% Trehaiose, pH 4.5.
RECONSTITUTION	Always centrifuge tubes before opening.Do not mix by vortex or pipetting.





It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at $< -20^{\circ}\text{C}$, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.

Aliquots of reconstituted samples are stable at $< -20^{\circ}\text{C}$ for 3 months.

BACKGROUND

Recombinant Human Interleukin-2 is a highly purified protein with a molecular weight of approximately 15,300 Daltons. The chemical name is des-alanyl-1, serine-145 Human Interleukin-2. It is produced by recombinant DNA technology using a genetically engineered E. coli strain containing an analog of the human interleukin-2 gene. Genetic engineering techniques were used to modify the Human IL-2 gene,





and the resulting expression clone encodes a modified Human IL-2. This recombinant form differs from native Interleukin-2 in following ways: it is not glycosylated; the molecule has serine substituted for cysteine at amino acid position 145; the aggregation state of molecule is likely to be different from that of native IL-2.

SDS-PAGE

