



# Recombinant Human Siglec-5 (C-6His-Flag-Fc)

<b>Catalog #</b>	EPT208
<b>Expression Host</b>	Human Cells
<b>DESCRIPTION</b>	Recombinant Human Sialic Acid-binding Ig-like Lectin 5 is produced by our Mammalian expression system and the target gene encoding Glu17-Thr434 is expressed with a 6His, Flag, Fc tag at the C-terminus.
<b>Accession</b>	O15389
<b>Synonyms</b>	Sialic acid-binding Ig-like lectin 5; Siglec-5; CD33 antigen-like 2; Obesity-binding protein 2; OB-BP2; CD170
<b>Mol Mass</b>	74.1 KDa
<b>AP Mol Mass</b>	90-110 KDa, reducing conditions
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Less than 0.1 ng/ $\mu$ g (1 EU/ $\mu$ g) as determined by LAL test.
<b>FORMULATION</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH





7.4.

## RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

## STORAGE

Lyophilized protein should be stored at  $< -20^{\circ}\text{C}$ , though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at  $4-7^{\circ}\text{C}$  for 2-7 days.

Aliquots of reconstituted samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.

## BACKGROUND

Human Siglec-5 are I type(Ig type) lectins belonging to the Ig superfamily, They are characterized by an N terminal Ig-like V type domain which mediates sialic acid binding, followed by varying numbers of Ig-like C2 type domains. SIGLEC5 has also been designated






CD170, they are expressed by monocytic or myeloid lineage cells, and also found at high levels in peripheral blood leukocytes, spleen, bone marrow and at lower levels in lymph node, lung, appendix, placenta, pancreas and thymus. SIGLEC5 are expressed by monocytes and neutrophils but absent from leukemic cell lines representing early stages of myelomonocytic differentiation. Siglec5 to 11 share a high degree of sequence similarity with CD33/Siglec3 both in their extracellular and intracellular regions. They are collectively referred to as CD33 related Siglecs. One remarkable feature of the CD33 related Siglecs is their differential expression pattern within the hematopoietic system. This fact, together with the presence of two conserved immunoreceptor tyrosinebased inhibition motifs (ITIMs) in their cytoplasmic tails, suggests that CD33 related Siglecs are involved in the regulation of cellular activation within the immune system.





**ELK Biotechnology**

## SDS-PAGE



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C